

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Prescription Solutions®

A UnitedHealth Group Company

PRIOR AUTHORIZATION CRITERIA – Formulary 2

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
Actemra	All FDA-approved indications not otherwise excluded from Part D.		<p>Rheumatoid Arthritis (RA):</p> <ol style="list-style-type: none"> 1. Diagnosis of moderate-to-severe RA; AND 2. Failed methotrexate or 2 DMARDs (azathioprine, cyclosporine, gold, hydroxychloroquine, leflunomide, penicillamine, sulfasalazine) for 3 months; AND 3. Failed 1 TNF antagonist <p>Reauthorization: Demonstration of clinical response to therapy.</p>	RA: 18 years and older.	RA (Initial): Prescribed or recommended by a rheumatologist	<p>Initial Therapy: 6 months</p> <p>Reauthorization: 12 months</p>	RA: Verification that the patient has been evaluated for tuberculosis and treated accordingly.
Adcirca	All FDA-approved indications not otherwise excluded from Part D.	Patients using organic nitrates.	<p>Pulmonary Arterial Hypertension: Patients with a confirmed diagnosis of pulmonary arterial hypertension (modified WHO Group I) which is symptomatic.</p>			Length of therapy	

Unison Advantage® (HMO) products are offered by UnitedHealthcare Insurance Company and its affiliated companies, Medicare Advantage organizations with a Medicare contract.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
Afinitor	All FDA-approved indications not otherwise excluded from Part D.					Length of the therapy	Failure of treatment with sunitinib or sorafenib.
Alimta	All FDA-approved indications not otherwise excluded from Part D.		<p>Malignant Pleural Mesothelioma</p> <ol style="list-style-type: none"> 1. Confirmed diagnosis of malignant pleural mesothelioma <p>Non-Small Cell Lung Cancer:</p> <ol style="list-style-type: none"> 1. Confirmed diagnosis of locally advanced or metastatic NSCLC; AND 2. One of the following: <ol style="list-style-type: none"> a. Prior history of first-line chemotherapy treatment for NSCLC. b. Used in combination with cisplatin. c. Tumor response or stable disease after 4 cycles of first-line platinum-based chemotherapy. d. Used as first-line monotherapy in patients 70 years of age or greater. e. Used as first-line monotherapy in patients less than 70 and not eligible for platinum-based chemotherapy. 			Length of the therapy	Alimta will be approved for continuation of prior therapy.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
Amevive	All FDA approved indications not otherwise excluded from Part D	CD4+ T-lymphocyte cell count < 250 cells per microliter.	<p>Initial Therapy for Plaque Psoriasis:</p> <ol style="list-style-type: none"> 1. Diagnosis of moderate-to-severe plaque psoriasis; AND 2. Failure to both phototherapy and systemic therapy with one of the following: methotrexate, cyclosporine, acitretin, hydroxyurea, sulfasalazine, 6-thioguanine, or mycophenolate. <p>Reauthorization for Plaque Psoriasis: Confirmation of positive clinical response to therapy</p>		Prescribed by a dermatologist.	<p>Initial Therapy: 12 weeks</p> <p>Re-authorization: 12 weeks</p>	Amevive will be reauthorized only if it has been at least 12 weeks since patient's last treatment
Ampyra	All FDA approved indications not otherwise excluded from Part D		<p>Initial Therapy for Multiple Sclerosis:</p> <ol style="list-style-type: none"> 1. Diagnosis of multiple sclerosis. 2. Physician confirmation that patient has difficulty walking. <p>Reauthorization for Multiple Sclerosis: Physician confirmation that the patient's walking improved with Ampyra therapy.</p>			<p>Initial Therapy: 3 months.</p> <p>Re-authorization: 1 year.</p>	
Anadrol-50	All FDA approved indications not otherwise excluded from Part D		<p>Acquired Aplastic Anemia:</p> <ol style="list-style-type: none"> 1. History of failure; OR 2. Used in combination with, antilymphocyte globulin or both antilymphocyte globulin and corticosteroid treatment. <p>Hypoplastic Anemia:</p> <ol style="list-style-type: none"> 1. Diagnosis of hypoplastic anemia due to myelotoxic drugs; AND 2. Failure to an erythropoietic stimulating agent. 			Length of the therapy for hypoplastic anemia. For other uses, authorization will be granted for 12 months.	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>Pure Red Cell Aplasia: Failure to immunosuppressive therapy.</p> <p>Chronic Renal Failure: Failure to an erythropoietic stimulating agent.</p>				
Apokyn	All FDA approved indications not otherwise excluded from Part D		<p>Advanced Parkinson's Disease:</p> <ol style="list-style-type: none"> 1. Confirmed diagnosis of advanced Parkinson's disease; AND 2. Unable to control "off" symptoms with adequate combinations of conventional oral therapy; AND 3. Used in combination with a non-5-HT3 antagonist antiemetic for initial therapy; AND 4. Not used in combination with 5-HT3 antagonists. 			1 year	Apokyn will only be approved for intermittent subcutaneous injection.
Aranesp	All FDA approved indications not otherwise excluded from Part D	<p>Anemia Due to Chronic Renal Failure: Patient is on dialysis (covered under Part B).</p> <p>Anemia in cancer patients on</p>	<p>Initial Therapy for Chronic Renal Failure:</p> <ol style="list-style-type: none"> 1. Hct < 33% OR Hgb < 11 gm/dl; AND 2. Verification of iron evaluation for adequate iron stores. <p>Reauthorization for Chronic Renal Failure:</p> <ol style="list-style-type: none"> 1. Verification that average Hct was below 36% over a 3-month period; AND 2. Verification of iron evaluation for adequate iron stores; AND 3. One of the following: <ol style="list-style-type: none"> a. Hct reached target range (30% to 36%); OR b. Decrease in blood transfusion; OR c. Hgb is ≥ 1 g/dL from pre-treatment level. 			<p>Initial Therapy: Three months for chemotherapy and MDS. Six months for CRF.</p> <p>Reauthorization: 12 months for CRF and MDS</p>	<p>Aranesp is subject to Part B vs. Part D review.</p> <p>Chemotherapy-Induced Anemia: Hgb/Hct levels must be collected within prior two weeks of</p>

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
		<p>chemotherapy: Patient is not receiving cancer chemotherapy or patient has malignancy for which therapy with Aranesp is contraindicated.</p> <p>Other off-label requests: Hgb greater than 10 gm/dL or Hct greater than 30%</p>	<p>Initial Therapy for Chemotherapy:</p> <ol style="list-style-type: none"> 1. Verification that other causes of anemia have been ruled out; AND 2. Verification of iron evaluation for adequate iron stores with one of the following: <ol style="list-style-type: none"> a. Hct < 36%; OR b. Hgb < 12 gm/dl; 3. Verification that the cancer is a non-myeloid malignancy. 4. AND one of the following: <ol style="list-style-type: none"> a. Verification that the patient is concurrently on chemotherapy; OR b. Will be on concomitant chemotherapy for 2 months; OR c. The anemia is caused by cancer chemotherapy. <p>Reauthorization for Chemotherapy:</p> <ol style="list-style-type: none"> 1. Hct < 36% OR Hgb < 12 gm/dl; AND 2. Hct reached target range (30% to 36%); AND 3. One of the following: <ol style="list-style-type: none"> a. Decrease in blood transfusion; OR b. Hgb is 1 g/dL; OR c. greater from pre-treatment level. 4. One of the following: <ol style="list-style-type: none"> a. Verification that the patient is concurrently on chemotherapy; OR b. Will be on concomitant chemotherapy for 2 months; OR c. The anemia is caused by cancer chemotherapy. <p>Initial Therapy for Myelodysplastic Syndrome:</p>				<p>request.</p> <p>All other uses: Hgb/Hct levels must be collected within prior 30 days of request.</p>

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			1. Hct < 33%; OR Hgb < 11 g/dL; AND 2. One of the following: a. Serum erythropoietin of ≤ 500 mU/mL; OR b. Diagnosis of transfusion-dependent MDS. 3. Verification of adequate iron stores. Reauthorization of Myelodysplastic Syndrome: 1. Verification that average Hct was below 36% over a 3 month period; AND 2. One of the following: a. Verification that Hct reached target (30% to 36%); OR b. Decrease in blood transfusion; OR c. Hgb increase ≥ 1 g/dL from pre-treatment level.				
Arcalyst	All FDA approved indications not otherwise excluded from Part D			12 years and older		Indefinite, long-term therapy (open-ended)	
Arzerra	All FDA approved indications not otherwise excluded from Part D		Chronic lymphocytic leukemia: 1. Confirmed diagnosis of chronic lymphocytic leukemia (CLL). 2. Refractory to both Campath and Fludara, or relapsed or refractory to two first-line CLL regimens containing Campath, Treanda, Leukeran, Cytosan, Fludara, Nipen, or Rituxan.			24 weeks	Prior authorization applies to new starts only
Avastin	All FDA	NSCLC:	Colorectal Cancer:		ARMD:	Colorectal	Avastin will be

Unison Advantage® (HMO) products are offered by UnitedHealthcare Insurance Company and its affiliated companies, Medicare Advantage organizations with a Medicare contract.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
	approved indications not otherwise excluded from Part D	<ol style="list-style-type: none"> 1. Squamous cell histology. 2. History of hemoptysis. 3. CNS metastases. 4. On-going therapeutic anticoagulation 	<ol style="list-style-type: none"> 1. Diagnosis of metastatic colorectal cancer; AND 2. Used in combination with: <ol style="list-style-type: none"> a. 5-FU; OR b. oxaliplatin plus capecitabine; OR c. capecitabine. <p>Non-Small Cell Lung Cancer:</p> <ol style="list-style-type: none"> 1. Diagnosis of unresectable locally advanced recurrent or metastatic NSCLC; AND 2. Used in combination with paclitaxel and carboplatin. <p>Renal Cell Cancer:</p> <ol style="list-style-type: none"> 1. Diagnosis of metastatic renal cell cancer; AND 2. Used in combination with interferon-alpha or refractory to either interferon alpha or interleukin-2. <p>Breast Cancer:</p> <ol style="list-style-type: none"> 1. Diagnosis of metastatic breast cancer; AND 2. Used in combination with paclitaxel. <p>Age-related Macular Degeneration::</p> <ol style="list-style-type: none"> 1. Failure to FDA-approved therapies or likely to have greater benefit from the use of intravitreal bevacizumab. 		Prescribed or recommended by retina specialist	Cancer, NSCLC, RCC, Breast Cancer, ARMD: Length of therapy	approved for continuation of prior therapy.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
Avonex	All FDA approved indications not otherwise excluded from Part D		Relapsing Multiple Sclerosis: 1. Recent history of a first clinical demyelinating event; AND 2. MRI-detected brain lesions consistent with MS.			1 year	
Berinert	All FDA approved indications not otherwise excluded from Part D		Hereditary Angioedema For the treatment of acute abdominal or facial Hereditary Angioedema attacks		Prescribed by an immunologist, allergist, or rheumatologist	12 months	
Betaseron	All FDA approved indications not otherwise excluded from Part D		Relapsing Multiple Sclerosis: 1. Patients with relapsing form of MS; OR 2. Patients with secondary progressive MS who continue to experience relapses.			1 year	
Botox	All FDA approved indications not otherwise excluded from Part D		Initial Therapy for Primary Axillary Hyperhidrosis: 1. History of failure to topical prescription strength drying agents; AND 2. One of the following: a. Score of 3 or 4 on the Hyperhidrosis Disease Severity Scale (HDSS); OR b. Skin maceration with secondary infection. Reauthorization for Hyperhidrosis: At least a 2-point improvement in HDSS.	VII cranial nerve disorders: ≥ 12 years of age	Initial therapy for Migraine : Prescribed by a neurologist or pain specialist. Initial Therapy for Chronic Back	Hyperhidrosis : One prescription Back Pain: One treatment (may be reauthorized for a maximum of two	

Unison Advantage® (HMO) products are offered by UnitedHealthcare Insurance Company and its affiliated companies, Medicare Advantage organizations with a Medicare contract.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>Initial Therapy for Migraine:</p> <ol style="list-style-type: none"> 1. Failure or contraindication to 5-HT1 agonists; AND 2. At least two prophylactic therapies for migraine; AND 3. Submission of chart documentation documenting complete evaluation of the patient. <p>Reauthorization for Migraine:</p> <ol style="list-style-type: none"> 1. Reduction in headache frequency; AND 2. Submission of chart notes documenting decreased utilization of pain medications or triptans, or a reduction in the number of emergency room visits. <p>Initial Therapy for Achalasia:</p> <ol style="list-style-type: none"> 1. High risk of complication from pneumatic dilation or surgical myotomy; OR 2. Failure to prior pneumatic dilation or surgical myotomy; OR 3. Prior dilation caused esophageal perforation; OR 4. Patient has an epiphrenic diverticulum or hiatal hernia. <p>Reauthorization for Achalasia:</p> <ol style="list-style-type: none"> 1. Documentation of improvement or reduction in symptoms of achalasia. <p>Initial Therapy for Anal Fissure:</p> <ol style="list-style-type: none"> 1. Failure of two conventional therapies. <p>Reauthorization for Anal Fissure:</p> <ol style="list-style-type: none"> 1. Incomplete healing of fissure or recurrence of fissure; AND 		<p>Pain: Prescribed by a neurologist, neurosurgeon, orthopedist, or pain specialist.</p> <p>Initial Therapy for Urinary Incontinence: Prescribed by a neurologist, neurosurgeon, or urologist.</p>	<p>treatments per year).</p> <p>Urinary Incontinence: 6 months for a single dose</p> <p>Other uses: 6 months</p>	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>2. Improved symptoms with prior treatment with Botox.</p> <p>Initial Therapy for Chronic Back Pain:</p> <ol style="list-style-type: none"> 1. Confirmed history of persistent low back pain > 6 months duration; AND 2. Failure to NSAIDs and opioids. <p>Reauthorization for Back Pain:</p> <p>Confirmed improvement in symptoms with initial Botox treatment.</p> <p>Initial Therapy for Urinary Incontinence:</p> <ol style="list-style-type: none"> 1. Failure to two oral anticholinergic agents; AND 2. One of the following conditions: <ol style="list-style-type: none"> a. Detrusor sphincter dyssynergia; OR b. Neurogenic detrusor overactivity <p>Reauthorization for Urinary Incontinence:</p> <ol style="list-style-type: none"> 1. Confirmed improvement in symptoms with initial Botox treatment; AND 2. At least 6 months has elapsed since the last Botox treatment. 				
Cellcept Intravenous	All FDA approved indications not otherwise excluded from Part D		<p>Transplant</p> <ol style="list-style-type: none"> 1. Patient received a renal (kidney), cardiac (heart), or hepatic (liver) transplant; AND 2. Patient is unable to take oral formulations of mycophenolate. <p>Lupus Nephritis:</p>			Length of therapy	Cellcept is subject to Part B vs. Part D review (not limited to new starts only).

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			1. Diagnosis of lupus nephritis; AND 2. Failure to combination therapy with corticosteroids and cyclophosphamide. 3. Patient is unable to take oral formulations of mycophenolate.				Cellcept will be approved for continuation of prior therapy if Part D.
Cellcept	All FDA approved indications not otherwise excluded from Part D		<p>Transplant:</p> 1. Patient received a renal (kidney), cardiac (heart), or hepatic (liver) transplant. 2. Patient received a bone marrow/stem cell transplant. <p>Lupus Nephritis:</p> 1. Diagnosis of lupus nephritis; AND 2. Failure to combination therapy with corticosteroids and cyclophosphamide. <p>Obliterative Bronchiolitis: Diagnosis of obliterative bronchiolitis following lung transplantation.</p>			Length of therapy	Cellcept is subject to Part B vs. Part D review (not limited to new starts only). Cellcept will be approved for continuation of prior therapy if Part D.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
Cesamet	All FDA approved indications not otherwise excluded from Part D		<p>Nausea and Vomiting Associated with Cancer Chemotherapy:</p> <ol style="list-style-type: none"> 1. Patient is receiving cancer chemotherapy 2. Failure to 5HT-3 receptor antagonist 3. Failure to one of the following agents: <ol style="list-style-type: none"> a. Antihistamine; OR b. Corticosteroid; OR c. Prokinetic agent; OR d. Antipsychotic 			Authorization will be issued for 6 months	<p>Cesamet is subject to Part B vs. Part D review.</p> <p>Cesamet will be approved for therapy continuation covered under Part B</p>
Chorionic Gonadotropin, Novarel, Pregnyl	All FDA approved indications not otherwise excluded from Part D					6 months	
Cimzia	All FDA approved indications not otherwise excluded from Part D		<p>Moderate to severe Crohn's disease: Initial Therapy:</p> <ol style="list-style-type: none"> 1 Failure to one of the following therapies supported by the American College of Gastroenterology Crohn's disease practice guideline: aminosalicylates, azathioprine, corticosteroids, methotrexate, or 6-mercaptopurine.; AND 2. Verification that patient has been evaluated for tuberculosis and treated accordingly. <p>Reauthorization: Demonstrated remission or significant clinical response to Cimzia therapy.</p>	18 years and older	<p>CD: Prescribed or recommended by gastroenterologist</p> <p>RA: Prescribed or recommended by a rheumatologist</p>	<p>CD Initial Therapy: 2 months</p> <p>CD Reauthorization: 12 months</p> <p>RA: 12 months</p>	

Unison Advantage® (HMO) products are offered by UnitedHealthcare Insurance Company and its affiliated companies, Medicare Advantage organizations with a Medicare contract.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>RA (initial therapy): 1. Diagnosis of moderately to severely active RA. 2. Failed MTX or 2 DMARDs for 3 mo. 3. Verification that the patient has been evaluated for tuberculosis and treated accordingly.</p> <p>Reauthorization: Submission of chart documentation demonstrating a positive clinical response.</p>				
Cinryze	All FDA approved indications not otherwise excluded from Part D		<p>Hereditary Angioedema 1. For the long term prevention of Hereditary Angioedema (HAE) attacks; AND 2. One of the following: a. Failure to danazol; OR b. Failure to one alkylated androgen</p>		Prescribed by an immunologist, allergist, or rheumatologist	12 months	
Copaxone	All FDA approved indications not otherwise excluded from Part D		<p>Relapsing Remitting Multiple Sclerosis: For patients with relapsing-remitting form of multiple sclerosis. Or for patients with history of a first clinical demyelinating event.</p>			1 year	
Degarelix	All FDA-approved indications not otherwise excluded from Part D.		Failure to an LHRH agonist			12 months	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
Emend	All FDA-approved indications not otherwise excluded from Part D.		<p>Acute Chemotherapy-Induced Nausea and Vomiting:</p> <ol style="list-style-type: none"> 1. Patient is currently receiving moderately or highly emetogenic chemotherapy; AND 2. Patient is concurrently on both a corticosteroid and a 5-HT3 receptor antagonist. <p>Delayed Chemotherapy-Induced Nausea and Vomiting:</p> <ol style="list-style-type: none"> 1. Patient is currently receiving highly emetogenic chemotherapy and a steroid; OR 2. Patient is on an anthracycline and cyclophosphamide. <p>Prevention of Postoperative Nausea and Vomiting:</p> <p>For the prevention of postoperative nausea and vomiting when administered prior to the induction of anesthesia.</p>			Acute CINV, Delayed CINV, PONV: 6 months	Emend is subject to Part B vs. Part D review.
Enbrel, Enbrel Sureclick	All FDA-approved indications not otherwise excluded from Part D.		<p>Rheumatoid Arthritis (RA):</p> <ol style="list-style-type: none"> 1. Diagnosis of moderate-to-severe RA; AND 2. Failed methotrexate or 2 DMARDs for 3 months. <p>Juvenile Idiopathic Arthritis (JIA):</p> <ol style="list-style-type: none"> 1. Diagnosis of moderate-to-severe poly-articular course JIA; AND 2. Failed NSAID or steroid and methotrexate for three 	<p>RA, PsA, AS, PPs: ≥ 18 years</p> <p>JIA: ≥ 2 years</p>	Initial Therapy for RA, JIA, PsA, AS: Prescribed or recommended by a rheumatolo-	Initial Therapy: 3 months for plaque psoriasis; 12 months for other uses	All diagnoses require verification that the patient has been evaluated for tuberculosis

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>months</p> <p>Psoriatic Arthritis (PsA): 1. Diagnosis of active PsA. 2. Failed methotrexate or 2 DMARDs for 3 months.</p> <p>Ankylosing Spondylitis (AS): 1. Diagnosis of AS. 2. Failed 2 NSAIDs for 3 months.</p> <p>Plaque Psoriasis (PPs): 1. Diagnosis moderate-to-severe chronic (greater than 6 months) plaque psoriasis. 2. Failed phototherapy and systemic therapy with one of the following: methotrexate, cyclosporine, acitretin, hydroxyurea, sulfasalazine, 6-thioguanine, or mycophenolate.</p> <p>Reauthorization: Demonstration of clinical response to therapy.</p>		<p>gist.</p> <p>Initial Therapy for PPs: Prescribed or recommended by a dermatologist.</p>	<p>Reauthorization: 12 months for all uses</p>	<p>and has been treated accordingly</p> <p>Reauthorization of Enbrel for PPs requires a dosage of 50 mg or less per week or less.</p>
Epogen, Procrit	All FDA-approved indications not otherwise excluded from Part D.	Anemia in cancer patients on chemotherapy: Patient is not receiving cancer chemo-	<p>Anemia due to Chronic Renal Failure: 1. Hematocrit (Hct) less than 33%; OR 2. Hemoglobin (Hgb) less than 11 gm/dl.</p> <p>Reauthorization of CRF: 1. Average Hct was below 36% over 3-months: AND 2. One of the following: a. Hct reached target (30% to 36%); OR b. Decrease in blood transfusion; OR c. Hgb is 1 g/dL or greater from pre-treatment level.</p>			<p>Initial Therapy Pre-Op: 1 month</p> <p>Chemo, HCV, and MDS: 3 months.</p> <p>CRF, HIV: 6 months</p>	<p>Epoetin will be subject to Part B vs. Part D review.</p> <p>For Chemotherapy-Induced Anemia, Hgb/Hct levels</p>

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
		therapy; OR Patient has malignancy for which therapy with epoetin is contra-indicated. Chronic Renal Failure: Patient is on dialysis (covered under Part B). Other off-label requests: Hgb greater than 10 gm/dL or Hct greater than 30%	Anemia in HIV-infected patients: 1. Anemia is due to zidovudine treatment or due to HIV infection; AND 2. Hgb less than 12 g/dL or Hct less than 36%. Reauthorization in HIV: 1. Hct was below 36% over 3 months; AND 2. One of the following: a. Hct reached target (30% to 36%); OR b. Decrease in blood transfusion; OR c. Hgb is 1 g/dL or greater from pre-treatment level. Anemia in cancer patients on Chemotherapy: 1. Verify other causes of anemia have been ruled out; AND 2. Hct less than 36% or Hgb less than 12 gm/dl. 3. Cancer is a non-myeloid malignancy; AND 4. Concurrently on chemo, will be on concomitant chemo for 2 months OR anemia is caused by cancer chemotherapy. Reauthorization in Chemo: 1. Hct less than 36% or Hgb less than 12 gm/dl; AND 2. One of the following; AND: a. Hct reached target (30% to 36%) b. Decrease in blood transfusion c. Hgb is 1 g/dL or greater from pre-treatment level. 3. Concurrently on chemotherapy for 2 months or anemia is caused by cancer chemo.			Reauthorization on CRF, HIV: 6 months HCV: 3 months. Other uses: 12 months	must be collected within prior two weeks of request. For all other indications, Hgb/Hct levels must be collected within prior 30 days of request.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>Preoperative use in patients undergoing surgery for reduction of allogeneic blood transfusion (Pre-op):</p> <ol style="list-style-type: none"> 1. Hgb greater than 10 to less than 13 g/dL scheduled to undergo elective, non-cardiac/vascular surgery to reduce blood transfusions; OR 2. Patient at high risk for perioperative transfusions with expected blood loss of 2 units or greater. <p>Refractory anemia in Myelodysplastic Syndrome:</p> <ol style="list-style-type: none"> 1. Hct less than 33% or Hgb less than 11 g/dL; AND 2. One of the following: <ol style="list-style-type: none"> a. Serum erythropoietin of 500 mU/mL or less b. Diagnosis of transfusion-dependent MDS. <p>Reauthorization for MDS:</p> <ol style="list-style-type: none"> 1. Average Hct was below 36% over a 3 months; AND 2. One of the following: <ol style="list-style-type: none"> a. Hct reached target (30% to 36%) b. Decrease in blood transfusion c. Hgb increase of 1 g/dL or more from pre- treatment level. <p>Treatment of anemia in HCV-infected patients due to ribavirin in combination with interferon or peg-interferon:</p> <ol style="list-style-type: none"> 1. Hgb less than 12 g/dL or Hct less than 36%; AND 2. Is concurrently on ribavirin and interferon or peg-interferon alfa for the treatment of HCV and the anemia is due to treatment. <p>Reauthorization of HCV:</p>				

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>1. Average Hct was below 36% over a 3 months; And</p> <p>2. One of the following:</p> <ul style="list-style-type: none"> a. Hct reached target (30% to 36%) b. Decrease in blood transfusion c. Hgb is 1 g/dL or greater from pre-treatment level. <p>All uses: Verify iron evaluation for adequate Fe stores.</p>				
Erbix	All FDA-approved indications not otherwise excluded from Part D.		<p>Head and Neck Cancer:</p> <ul style="list-style-type: none"> 1. One of the following: <ul style="list-style-type: none"> a. Confirmed diagnosis of locally or regionally advanced squamous cell carcinoma of the head and neck b. Recurrent or metastatic squamous cell head and neck cancer; AND 2. One of the following: <ul style="list-style-type: none"> a. Used in combination with radiation therapy b. After failure of platinum-based chemotherapy. <p>Colorectal Cancer:</p> <ul style="list-style-type: none"> 1. Confirmed diagnosis of metastatic carcinoma of the colon or rectum; AND 2. One of the following: <ul style="list-style-type: none"> a. Used in combination with irinotecan-based chemotherapy b. Intolerance to irinotecan-based chemotherapy c. Failure of irinotecan or oxaliplatin-based chemotherapy regimens; AND 3. Tumor expresses wild-type KRAS gene. 			Length of therapy	Erbix will be approved for continuation of prior therapy.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
Extavia	All FDA-approved indications not otherwise excluded from Part D.					12 months	
fentanyl citrate oral trans-mucosal, Fentora, Onsolis	All FDA-approved indications not otherwise excluded from Part D.		Cancer Pain: 1. Confirmed diagnosis of malignant pain; AND 2. Failure or contraindication to an immediate-release opioid; AND 3. Demonstrated tolerance to opioids.			Length of therapy	
Folotyn	All FDA-approved indications not otherwise excluded from Part D.		Peripheral T-cell lymphoma 1. Diagnosis of relapsed or refractory peripheral T-cell lymphoma 2. Verification that patient is receiving folic acid and vitamin B12 supplementation.			12 months	Folotyn will be approved for continuation of prior therapy.
Forteo	All FDA-approved indications not otherwise excluded from Part D.	1. Paget's disease history 2. Bone metastases of skeletal malignancies	Osteoporosis: BMD T score of -3.0 or less and a previous fracture resulting from minimal trauma, or both of the following: 1. Failure to a formulary bisphosphonate; AND 2. Patient has a history of fracture resulting from minimal trauma or BMD T score of -2.5 or less.			2 years	Forteo is subject to Part B vs. Part D review.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
		3. Radiation therapy 4. Bone disease other than osteoporosis 5. Concurrent use of bisphosphonate.					
Gamastan S/D	All FDA approved indications not otherwise excluded from Part D		<p>Hepatitis A: For use before or soon after exposure.</p> <p>Measles: For use in susceptible individuals exposed fewer than 6 days previously.</p> <p>Varicella: For use in immunocompromised patients.</p> <p>Rubella: For pregnant women who will not consider a therapeutic abortion.</p>			Length of therapy	
Gleevec	All FDA approved indications not		<p>Chronic Myeloid Leukemia (Adults): Diagnosis of Philadelphia chromosome positive CML.</p>			Length of therapy	Gleevec will be approved for

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
	otherwise excluded from Part D		<p>Chronic Myeloid Leukemia (Children): 1. Diagnosis of Philadelphia chromosome positive (Ph+) chronic phase CML; AND 2. One of the following: a. Not candidates for stem cell transplantation b. Disease has recurred after stem cell transplant c. Patients who are resistant to interferon-alfa therapy.</p> <p>Acute Lymphoblastic Leukemia: Adult patients with Philadelphia chromosome positive ALL.</p> <p>Myelodysplastic/Myeloproliferative disease: Adults diagnosed with MDS/MPD diseases associated with platelet-derived growth factor receptor gene rearrangements.</p> <p>Aggressive systemic mastocytosis: 1. Adults diagnosed with aggressive systemic mastocytosis; AND 2. One of the following: a. Patient is without the D816V c-Kit mutation b. c-Kit mutation status unknown.</p> <p>Hypereosinophilic syndrome and chronic eosinophilic leukemia : Adults diagnosed with HES or CEL.</p> <p>Dermatofibrosarcoma protuberans: Adults with unresectable, recurrent and/or metastatic DFSP.</p>				continuation of prior therapy.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>Gastrointestinal Stromal Tumors: Patients with a confirmed diagnosis of unresectable and/or metastatic GIST.</p>				
<p>Growth Hormones: Genotropin, Genotropin Miniquick, Humatrope, Humatrope Combo Pack, Norditropin Cartridge, Norditropin Nordiflex Pen, Nutropin, Nutropin Aq, Nutropin Aq Pen, Omnitrope, Saizen, Saizen Click.Easy, Tev-Tropin</p>	<p>All FDA approved indications not otherwise excluded from Part D:</p> <p>Growth Hormone Deficiency (GHD) in Children</p> <p>Prader-Willi Syndrome (PWS)</p> <p>Small for Gestational Age (SGA)</p> <p>Turner Syndrome (TS)</p> <p>Noonan Syndrome (NS)</p>	<p>Chidhood Onset Growth Hormone Deficiency in Adults:</p> <ol style="list-style-type: none"> 1. Males with bone age greater than 17 yrs or females with bone age greater than 15 years 2. Closed bone epiphyses on radiograph 3. Growth velocity less than 	<p>GHD Children:</p> <ol style="list-style-type: none"> 1. Diagnosis of GH deficiency based on two GH stimulation tests or low Insulin-like growth factor 1 (IGF-1) levels; AND 2. Demonstrate growth failure based on growth velocity or height shorter than 2 standard deviations (SD) below the mean height for age. <p>Prader-Willi Syndrome or Small for Gestational Age:</p> <ol style="list-style-type: none"> 1. Diagnosis of PWS confirmed by genetic testing; OR 2. Diagnosis of SGA confirmed by birth wt of less than 2500g at gestation of more than 37 wks or at birth weight or length below the 3rd percentile for gestational age who failed to catch up by 2 years of age. <p>Turner Syndrome, Noonan Syndrome:</p> <ol style="list-style-type: none"> 1. Treatment of short stature in females w/bone age less than 15 years associated w/TS or NS; OR 2. Treatment of short stature in males w/bone age less than 17 years associated w/NS. <p>Growth Retardation associated with Chronic Renal Insufficiency:</p> <ol style="list-style-type: none"> 1. Diagnosis of chronic renal insufficiency; AND 2. Height shorter than or equal to 2 SD below the median 		<p>GHD (Child), AOGH, COGHDA, IGHDA, Initial Therapy for TS or NS, GRCRF, and ISS: Prescribed by an endocrinologist.</p>	<p>Length of therapy for GHD in adults. One year for all other uses.</p>	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
	<p>Growth Retardation associated with Chronic Renal Insufficiency (GRCRF)</p> <p>Idiopathic Short Stature (ISS)</p> <p>Adult Onset Growth Hormone Deficiency (AOGHD),</p> <p>Childhood Onset GH Deficiency in Adults (COGHDA)</p> <p>Isolated GH Deficiency in Adults (IGHDA)</p>	<p>2 cm/year during previous year of treatment unless COGHD criteria are met.</p>	<p>age for children or where growth velocity falls to below 4.5 cm/year.</p> <p>Reauthorization for GHD in Children, PWS, SGA, TS, NS, GRCRF:</p> <ol style="list-style-type: none"> 1. Increase in growth velocity of at least 2 cm/year during previous year of treatment; AND 2. Males with bone age less than 17 yrs or females with bone age less than 15 years. <p>Idiopathic Short Stature:</p> <ol style="list-style-type: none"> 1. Height less than or equal to 2.25 SD below the mean height for age. Growth velocity less than the 25th percentile for bone age; AND 2. Verify open epiphyses on last bone age radiograph; AND 3. Absence of comorbid conditions that should be observed or treated by other means. <p>Reauthorization of ISS:</p> <ol style="list-style-type: none"> 1. Increase in growth velocity of at least 4.5 cm/year during previous year of treatment; AND 2. Males w/bone age less than 17 years or females w/bone age less than 15 years. <p>Adult Onset Growth Hormone Deficiency:</p> <ol style="list-style-type: none"> 1. Pts who have GHD alone or multiple hormone deficiencies because of pituitary disease/insult, hypothalamic disease, surgery, or radiation treatment; AND 2. IGF-1 level less than 77 mcg/L or 2 SD below the 				

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>mean value, matched by age and gender.</p> <p>Childhood Onset GH Deficiency in Adults: 1. Childhood onset in patients who were GH deficient during childhood who have GH deficiency confirmed as an adult before replacement treatment with GH is started; AND 2. Persistent deficiency of GH documented by GH stimulation tests.</p> <p>Isolated GH Deficiency in Adults: Documented deficiency of GH documented by 2 GH stimulation tests.</p>				
Hexalen	All FDA approved indications not otherwise excluded from Part D.		<ol style="list-style-type: none"> 1. Diagnosis of ovarian cancer 2. Cancer has progressed or recurred following first-line treatment with a cisplatin or alkylating agent-based combination 		Prescribed by an oncologist	12 months	Hexalen will be approved for continuation of prior therapy.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
Humira, Humira Pen-Crohns Disease	All FDA approved indications not otherwise excluded from Part D.		<p>Rheumatoid Arthritis: 1. Diagnosis of moderate-to-severe RA; AND 2. Failed methotrexate or 2 DMARDs for 3 months.</p> <p>Juvenile Rheumatoid Arthritis / Juvenile Idiopathic Arthritis: Diagnosis of moderate-to-severe poly-articular course</p> <p>Juvenile Rheumatoid Arthritis: Failed NSAID or steroid and DMARD for three months.</p> <p>Psoriatic Arthritis: 1. Diagnosis of active PsA. 2. Failed methotrexate or 2 DMARDs for 3 months.</p> <p>Ankylosing Spondylitis: 1. Diagnosis of AS. 2. Failed 2 NSAIDs for 3 months.</p> <p>Plaque Psoriasis: 1. Diagnosis moderate-to-severe plaque psoriasis. 2. Failed phototherapy and systemic therapy.</p> <p>Crohn's disease: 1. Diagnosis of moderate to severe CD; AND 2. Failed conventional therapies.</p> <p>Reauthorization: Demonstration of clinical response to therapy.</p>	<p>RA, PsA, CD, AS, Plaque Psoriasis: 18 years and older.</p> <p>JIA: 4 years and older.</p>	<p>RA, PsA, AS, JIA: Prescribed or recommended by a rheumatologist.</p> <p>Plaque Psoriasis: Prescribed or recommended by a dermatologist.</p> <p>CD: Prescribed or recommended by gastroenterologist.</p>	<p>Initial Authorization: 4 months for Plaque Psoriasis; 12 months for other uses.</p> <p>Reauthorization 12 months</p>	<p>RA: Authorization is for 40 mg every other week unless documented treatment failure to Humira every other week dosing. Then Humira may be approved for every week dosing if other criteria met.</p> <p>Plaque Psoriasis: Humira dosage is 40 mg every other week.</p> <p>All diagnoses: Verification that the patient has been evaluated for TB and treated accordingly.</p> <p style="text-align: right;">10/1/10</p>

Unison Advantage® (HMO) products are offered by UnitedHealthcare Insurance Company and its affiliated companies, Medicare Advantage organizations with a Medicare contract.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
<p>Immune Globulin: Carimune Nanofiltered , Flebogamma, Gammagard Liquid, Gamunex, Octagam, Polygam S/D, Vivaglobin</p>	<p>All FDA approved indications not otherwise excluded from Part D</p>		<p>Idiopathic Thrombocytopenic Purpura (ITP): For patients with ITP who require a rapid temporary increase in platelet count or to control excessive bleeding</p> <p>Kawasaki Disease (KD): Confirmed diagnosis of KD.</p> <p>B-cell Chronic Lymphocytic Leukemia (CLL): 1. Documented hypogammaglobulinemia (IgG less than 600mg/dL); OR 2. History of bacterial infections associated with B-cell CLL.</p> <p>Bone Marrow Transplantation (BMT): 1. Confirmed allogeneic BMT within the last 100 days; AND 2. Documented severe hypogammaglobulinemia (IgG less than 400 mg/dL)</p> <p>Dermatomyositis and Polymyositis: Failure or intolerance to one of the following: corticosteroid therapy, methotrexate, azathioprine, or cyclophosphamide.</p> <p>HIV: Documented hypogammaglobulinemia (IgG less than 400 mg/dL).</p> <p>Guillane-Barre Syndrome (GBS): 1. Confirmed diagnosis of severe GBS; AND 2. Patients with severe disease requiring aid to walk;</p>		<p>MG: Prescribed by a neurologist.</p>	<p>BMT: 100 days after transplant</p> <p>KD: 1 month</p> <p>MG, GBS: 1 treatment course</p> <p>ITP, LEMS: 6 months</p> <p>Other Uses: 1year</p>	<p>Immune Globulin is subject to Part B vs. Part D review.</p> <p>For Part D: For patients in which immune globulin is administered in the patient's home.</p>

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>AND</p> <p>3. Onset of muscle weakness within the last 4 weeks.</p> <p>Lambert-Eaton Myasthenic Syndrome (LEMS):</p> <p>1. Confirmed diagnosis of LEMS.</p> <p>Acute Myasthenia Gravis (MG) Exacerbation :</p> <p>1. Confirmed diagnosis of myasthenia gravis with myasthenic exacerbation, defined by one of the following:</p> <ul style="list-style-type: none"> a. Difficulty swallowing b. Acute respiratory failure c. Major functional disability responsible for the discontinuation of physical activity. <p>Relapsing-Remitting Multiple Sclerosis (MS) :</p> <p>1. Confirmed diagnosis of relapsing remitting form of MS</p> <p>AND</p> <p>2. Failure to two of the following: Betaseron, Avonex, Rebif, Copaxone, Tysabri.</p> <p>Stiff Person Syndrome :</p> <p>Chart documentation confirming a diagnosis of stiff-person syndrome.</p>				
Infergen	All FDA approved indications not otherwise excluded from Part D		<p>Hepatitis C - Treatment Naive Patients:</p> <p>For patients with Chronic Hepatitis C with compensated liver disease with positive HCV antibody and HCV RNA.</p> <p>Hepatitis C - Continuation of Therapy:</p> <p>For genotypes 2, 3, 5, or 6: Loss of detectable HCV</p>	Hepatitis C in Treatment Naive patients:		Treatment-Naive: genotypes 2, 3, 5, 6: 6 months	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			RNA from serum or 100 fold drop or more in HCV RNA level.	18 years and older.		Genotypes 1, 4 or HIV/HCV: 12 months Continuation of treatment in genotypes 2, 3, 5, 6: 6 months	
Insulin-like Growth Factor: Increlex	All FDA approved indications not otherwise excluded from Part D		IGF-1 deficiency 1. All of the following: a. Diagnosis of severe primary IGF-1 deficiency. b. Height standard deviation score of -3.0 or less. c. Basal IGF-1 standard deviation score of -3.0 or less. d. Normal or elevated growth hormone. e. Open finger epiphyses on last bone radiograph GH gene deletion: a. Diagnosis of growth hormone gene deletion who have developed neutralizing antibodies to GH; AND b. Have open finger epiphyses on last bone radiograph.			All Uses: Insulin-like growth factor will be approved for 1 year.	
Intron-A, Intron-A W/Diluent	All FDA approved indications not otherwise excluded from Part D		Hepatitis B - HBeAg positive: 1. HBsAg positive for at least 6 months; AND 2. HBV DNA level greater than 100,000 copies/mL; AND. 3. Compensated liver disease; AND 4. One of the following: persistent ALT 2 times ULN or moderate to severe hepatitis or fibrosis on biopsy.	Hep B - HBeAg positive, Hep B - HBeAg negative :		HepB+: 6 months HepB-: 1 year HepC:	Intron A will be approved for continuation of prior therapy for neoplastic

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>Hepatitis B - HBeAg negative: 1. HBsAg positive for at least 6 months; AND 2. HBV DNA level of 2000 IU/mL or more or 11,200 copies/mL; AND 3. Compensated liver disease; AND 4. One of the following: persistent ALT 2 times ULN or moderate to severe hepatitis or fibrosis on biopsy.</p> <p>Hepatitis C - Treatment Naive Patients: For patients with Chronic Hepatitis C with compensated liver disease with positive HCV antibody and HCV RNA.</p> <p>Hepatitis C - Continuation of Therapy: For genotypes 2, 3, 5, or 6: loss of detectable HCV RNA from serum or 100 fold drop or more in HCV RNA level.</p> <p>Non-Hepatitis Diagnoses Diagnosis of one of the following: 1. Malignant Melanoma 2. Hairy cell leukemia (HCL) 3. Stage III or IV follicular Non-Hodgkin's Lymphoma 4. Condylomata acuminata 5. AIDS-related Kaposi's sarcoma 6. Multiple Myeloma.</p> <p>Acute Hepatitis C: Patients with acute hepatitis C.</p>	<p>1 year of age or older.</p> <p>Hep C - Treatment Naive Patients, Non-Hepatitis Diagnoses, Acute Hep C: 18 years old and older.</p> <p>Hep C - Treatment Naive Patients (in combination with ribavirin) : 3 years of age and older.</p>		<p>genotypes 2, 3, 5, 6: 6 months</p> <p>HepC: genotypes 1, 4, HIV/HCV: 12 months</p> <p>Acute HepC, HCL, Kaposi: 6 months.</p> <p>Warts: 3 weeks.</p> <p>Other uses: 1year</p>	<p>diseases.</p>

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
Istodax	All FDA approved indications not otherwise excluded from Part D		Cutaneous T-Cell Lymphoma (CTCL): Confirmed diagnosis of CTCL. Failure to one systemic therapy.			12 months	Istodax will be approved for continuation of prior therapy.
Kalbitor	All FDA approved indications not otherwise excluded from Part D		Hereditary Angioedema For the treatment of acute Hereditary Angioedema attack.		Prescribed by an immunologist, allergist, or rheumatologist	12 months	
Ketek	All FDA approved indications not otherwise excluded from Part D		Community-Acquired Pneumonia: 1. Diagnosis of CAP in an adult outpatient; AND 2. Resistance or failure to either azithromycin or clarithromycin.			Ketek will be approved for the length of therapy.	
Kineret	All FDA approved indications not otherwise excluded from Part D		Initial Therapy for Rheumatoid Arthritis (RA): 1. Moderate to severe active RA; AND 2. Failure with a TNF-alpha-blocker; AND 3. Failure on either methotrexate or at least 1 DMARD for at least 3 months. Reauthorization for RA: Submission of chart documentation demonstrating positive clinical response.	RA: 18 years and older	RA: Prescribed or recommended by a rheumatologist.	Initial Therapy for RA: Kineret will be approved for 12 months. Reauthorization for RA: Kineret will be approved for 1	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
						year.	
Letairis	All FDA approved indications not otherwise excluded from Part D		Pulmonary Arterial Hypertension: Patients with a confirmed diagnosis of pulmonary arterial hypertension (modified WHO Group I) which is symptomatic.			Length of therapy	
Leukine	All FDA approved indications not otherwise excluded from Part D		Bone Marrow/Stem Cell Transplant (BMSCT): 1. Patients with non-myeloid malignancies undergoing myeloablative chemotherapy followed by autologous/allogeneic BMT; OR 2. Mobilization of hematopoietic progenitor cells into the peripheral blood for collection by leukapheresis; OR 3. Peripheral stem cell transplant patients who have received myeloablative chemotherapy. Acute Myeloid Leukemia (AML) Induction or Consolidation Therapy: For patients with AML following induction or consolidation chemotherapy. Neutropenia associated with dose dense chemotherapy (NDDC): 1. Patient is receiving the National Comprehensive Cancer Network's (NCCN's) Breast Intergroup, INT C9741 dose dense chemotherapy protocol for primary breast cancer; OR 2. A dose-dense regimen for which the incidence of febrile neutropenia is unknown.	AML: Greater than or equal to 55 years old.		BMSCT, AML, NDDC, CFN, FN: 3 months HIVN: 6 months	

Unison Advantage® (HMO) products are offered by UnitedHealthcare Insurance Company and its affiliated companies, Medicare Advantage organizations with a Medicare contract.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>Chemotherapy with risk of febrile neutropenia (CFN): 1. Patient is receiving a chemotherapy regimen associated with more than 20% incidence of febrile neutropenia; OR 2. Patient is receiving chemotherapy regimen associated with 10-20% incidence of febrile neutropenia and has risk factors associated with chemotherapy-induced infection, febrile neutropenia or neutropenia.</p> <p>Febrile Neutropenia (FN): 1. For patients receiving myelosuppressive anticancer drugs associated with neutropenia; AND 2. Patient either has febrile neutropenia or has a history of febrile neutropenia during a previous course of chemotherapy.</p> <p>HIV-related neutropenia (HIVN): HIV-infected patients with an Absolute Neutrophil Count (ANC) less than or equal to 1,000 cells/mm³ with or without one or more risk factors for developing chronic neutropenia.</p>				
Lotronex	All FDA approved indications not otherwise excluded from Part D	Initial therapy for Irritable Bowel Syndrome (IBS) in the male gender.	<p>Initial Therapy for Irritable Bowel Syndrome: 1. Confirmed diagnosis of IBS with diarrhea predominant symptoms for at least 6 months; AND 2. Failure to an antispasmodic and an anti-diarrhea agent.</p> <p>Reauthorization for IBS: 1. Recurrence of diarrhea-predominant IBS; AND</p>	18 years and older.	Verification that physician has enrolled in the GlaxoSmith-Kline Prescribing Program.	<p>Initial Therapy: 12 weeks</p> <p>Reauthorization: 6 months</p>	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			2. Documentation of positive clinical response while on Lotronex.				
dronabinol	All FDA approved indications not otherwise excluded from Part D		<p>Nausea and Vomiting Associated with Cancer Chemotherapy (CINV):</p> <ol style="list-style-type: none"> 1. Patient is receiving cancer chemotherapy; AND 2. Failure to 5HT-3 receptor antagonist; AND 3. Failure to one of the following agents: <ol style="list-style-type: none"> a. Antihistamine b. Corticosteroid c. Prokinetic agent d. Antipsychotic. <p>AIDS Anorexia: Diagnosis of anorexia with weight loss in patients with AIDS.</p>			<p>CINV: 6 months</p> <p>AIDS anorexia: Length of therapy</p>	<p>Marinol is subject to Part B vs. Part D review.</p> <p>CINV: Marinol will be approved for continuation covered under Part B when patient is receiving chemotherapy .</p>
Meprobamate	All FDA approved indications not otherwise excluded from Part D			No prior authorization if patient is less than 65 years old		4 months	
Miacalcin injectable	All FDA approved indications not otherwise excluded from Part D		<p>Postmenopausal Osteoporosis:</p> <ol style="list-style-type: none"> 1. Failure to a bisphosphonate or selective estrogen-receptor modulator (SERM); AND 2. Failure to Miacalcin Nasal Spray; AND 3. History of vertebral compression fractures, or fractures of the hip or distal radius resulting from minimal 			<p>Post-menopausal Osteoporosis: 12 months</p> <p>Hypercalcemi</p>	Miacalcin is subject to Part B vs. Part D review.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			trauma, or T score of -2.5 or less. Initial Therapy for Paget's Disease: History of failure or intolerance to oral bisphosphonates. Reauthorization for Paget's Disease: Serum alkaline phosphatase concentration fails to normalize after the previous 6 months of therapy. Hypercalcemia: 1. Corrected total serum calcium of 12 mg/dl; OR 2. Greater or corrected total serum calcium of 6 mEq/L or greater.			a: 1 month Paget's Disease: 6 months	
Mozobil	All FDA-approved indications not otherwise excluded from Part D.		Used in combination with granulocyte-colony stimulating factor (G-CSF).		Prescribed by a hematologist/ oncologist	One course of therapy up to 4 days	
Multaq	All FDA approved indications not otherwise excluded from Part D	1. Patients with NYHA Class IV heart failure or NYHA Class II - III heart failure	Failure, intolerance, or contraindication to amiodarone.			1 year	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
		with a recent decompensation requiring hospitalization or referral to a specialized heart failure clinic. 2. Second- or third-degree atrioventricular (AV) block or sick sinus syndrome (except when used in conjunction with a functioni					

Unison Advantage® (HMO) products are offered by UnitedHealthcare Insurance Company and its affiliated companies, Medicare Advantage organizations with a Medicare contract.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
		ng pacemak -er). 3. Bradycar -dia less than 50 bpm. 4. QTc interval greater than or equal to 500 ms. 5. Severe hepatic impair- ment.					
Neulasta	All FDA approved indications not otherwise excluded from Part D		Chemotherapy with risk of febrile neutropenia (CFN): 1. Patient is receiving a chemotherapy regimen associated with more than 20% incidence of febrile neutropenia; OR 2. a. Patients is receiving chemotherapy regimen associated with 10-20% incidence of febrile neutropenia; AND b. Has risk factors associated with chemotherapy-induced infection, febrile neutropenia or neutropenia. Neutropenia associated with dose dense chemotherapy (NDDC):			CFN, NDDC, FN: One month or duration of treatment.	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>1. Patients is receiving NCCN's Breast Intergroup, INT C9741 dose dense chemotherapy protocol for primary breast cancer; OR</p> <p>2. A dose-dense regimen for which the incidence of febrile neutropenia is unknown.</p> <p>Febrile Neutropenia (FN):</p> <p>1. For patients receiving myelosuppressive anticancer drugs associated with neutropenia; AND</p> <p>2. Patient either has febrile neutropenia or has a history of febrile neutropenia during a previous course of chemotherapy.</p>				
Neumega	All FDA approved indications not otherwise excluded from Part D	Patients with myeloablative chemotherapy.	<p>Thrombocytopenia following chemotherapy</p> <p>1. Verification that the cancer is a non-myeloid malignancy; AND</p> <p>2. Platelet count is less than 50,000 cells/microliter; AND</p> <p>3. Patients with one or more of the following risk factors:</p> <ul style="list-style-type: none"> a. Extensive prior cytotoxic chemotherapy b. Prior severe chemotherapy-induced thrombocytopenia c. Receiving chemotherapy regimens associated with high risk for thrombocytopenia. 			3 week intervals for up to 6 cycles post-chemotherapy.	
Neupogen	All FDA approved indications not otherwise excluded from Part D		<p>Bone Marrow/Stem Cell Transplant (BMSCT):</p> <p>1. For patients with non-myeloid malignancies undergoing myelo-ablative chemotherapy followed by autologous or allogeneic BMT; OR</p> <p>2. For mobilization of hematopoietic progenitor cells into the peripheral blood for collection by leukapheresis; OR</p>			<p>BMSCT, AML, NDDC, CFN, FN: 3 months</p> <p>SCN, HCN: 12 months</p>	

Unison Advantage® (HMO) products are offered by UnitedHealthcare Insurance Company and its affiliated companies, Medicare Advantage organizations with a Medicare contract.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>3. For peripheral stem cell transplant patients who have received myelo-ablative chemotherapy.</p> <p>Acute Myeloid Leukemia (AML) Induction or Consolidation Therapy: For patients with AML following induction or consolidation chemotherapy.</p> <p>Neutropenia associated with dose dense chemotherapy (NDDC):</p> <ol style="list-style-type: none"> 1. Patient is receiving NCCN's Breast Intergroup, INT C9741 dose dense chemotherapy protocol for primary breast cancer; OR 2. A dose-dense regimen for which the incidence of febrile neutropenia is unknown. <p>Chemotherapy with risk of febrile neutropenia (CFN):</p> <ol style="list-style-type: none"> 1. Patient is receiving a chemotherapy regimen associated with more than 20% incidence of febrile neutropenia; OR 2. Patient is receiving a chemotherapy regimen associated with 10-20% incidence of febrile neutropenia and has risk factors associated with chemotherapy-induced infection, febrile neutropenia or neutropenia. <p>Febrile Neutropenia (FN):</p> <ol style="list-style-type: none"> 1. For patients receiving myelosuppressive anticancer drugs associated with neutropenia; AND 2. Patient either has febrile neutropenia or has a history of febrile neutropenia during a previous course of 			<p>HIVN: 6 months</p>	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>chemotherapy.</p> <p>Severe Chronic Neutropenia (SCN): For patients with severe chronic neutropenia.</p> <p>Hepatitis-C Treatment of Related Neutropenia (HCN): 1. Neutropenia in Hepatitis C virus infected patients undergoing treatment with Peg-Intron or Pegasys after dose reduction; OR 2. For patients with HIV co-infection or status post liver transplant, or established cirrhosis who experience interferon-induced neutropenia due to treatment with Peg-Intron or Pegasys.</p> <p>HIV-related neutropenia (HVN): HIV-infected patients with an ANC less than or equal to 1,000 cells/mm³ with or without one or more risk factors for developing chronic neutropenia.</p>				
Nexavar	All FDA approved indications not otherwise excluded from Part D		<p>One of the following:</p> <ol style="list-style-type: none"> 1. Diagnosis of renal cell carcinoma with relapse following surgical excision 2. Diagnosis of renal cell carcinoma with medically or surgically unresectable tumor 3. Diagnosis of Stage IV renal cell carcinoma 4. Diagnosis of unresectable hepatocellular carcinoma. 		Prescribed by an oncologist.	6 months	Nexavar will be approved for continuation of prior therapy.
Opium tincture	All FDA approved indications not otherwise		<p>Initial Therapy for Diarrhea:</p> <ol style="list-style-type: none"> 1. Treatment of diarrhea 2. Failed, contraindication, or intolerance to Imodium (loperamide) and Lomotil (diphenoxylate/atropine). 	18 years and older.	Prescribed by a gastroenterologist or in	1 year	

Unison Advantage® (HMO) products are offered by UnitedHealthcare Insurance Company and its affiliated companies, Medicare Advantage organizations with a Medicare contract.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
	excluded from Part D				consultation with a gastroenterologist.		
Orencia	All FDA approved indications not otherwise excluded from Part D		<p>Rheumatoid Arthritis (RA): 1. Diagnosis of moderate-to-severe RA; AND 2. Failed methotrexate or 2 DMARDs for 3 months.</p> <p>Reauthorization for RA: Documentation of clinical improvement from ongoing therapy.</p> <p>Juvenile Idiopathic Arthritis (JIA): Failed NSAID or steroid and methotrexate for three months.</p> <p>Reauthorization for JIA: Documentation of clinical improvement from ongoing therapy.</p>	<p>RA: 18 years and older.</p> <p>JIA: 6 years and older.</p>	Prescribed or recommended by a rheumatologist.	12 months.	Verification that the patient has been evaluated for tuberculosis and treated accordingly.
oxandrolone	All FDA approved indications not otherwise excluded from Part D		<p>Bone Pain: Diagnosis of bone pain due to osteoporosis.</p> <p>Initial Therapy for AIDS Wasting: Diagnosis of AIDS wasting/cachexia and failure to hormone replacement therapy in patients with hypogonadism.</p> <p>Reauthorization for AIDS Wasting: Verification that the patient's weight has increased a</p>			<p>Initial therapy: 3 months</p> <p>Reauthorization: Length of therapy</p>	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			minimum of 2% while taking Oxandrin				
Pegasys	All FDA approved indications not otherwise excluded from Part D		<p>Hepatitis B - HBeAg positive patients:</p> <ol style="list-style-type: none"> 1. HBsAg positive for at least 6 months; AND 2. HBV DNA level greater than 100,000 copies/mL; AND. 3. Compensated liver disease; AND 4. One of the following: <ol style="list-style-type: none"> a. ALT (liver enzyme) 2 times upper limits of normal (ULN) b. Moderate-to-severe hepatitis or fibrosis on biopsy. <p>Hepatitis - HBeAg negative patients:</p> <ol style="list-style-type: none"> 1. HBsAg positive for at least 6 months; AND 2. HBV DNA level of 2000 IU/mL or more or 11,200 copies/mL; AND 3. Compensated liver disease; AND 4. One of the following: <ol style="list-style-type: none"> a. ALT 2 times ULN b. Moderate-to-severe hepatitis or fibrosis on biopsy. <p>Hepatitis C - Treatment Naive Patients:</p> <ol style="list-style-type: none"> 1. Chronic Hepatitis C with compensated liver disease; AND 2. Positive HCV antibody HCV RNA; AND 3. HCV RNA level measurement; AND 4. Genotype test result; AND 5. For patients who have not previously been treated with interferon. <p>Continuation of Therapy:</p> <p>A. For genotypes 5 or 6:</p>	For all covered uses: 18 years and older		<p>Hepatitis B: 1year.</p> <p>Hepatitis C Genotypes 5, 6: 12 weeks; Genotypes 2, 3: 24 weeks; Genotypes 1, 4: (HIV/HCV co-infected patients): 48wk.</p> <p>Hepatitis C Continuation therapy: Genotypes 1,3: 24 weeks, Genotypes 5, 6: 36wk.</p> <p>Hepatitis C Retreatment: 1year</p>	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>1. Loss of detectable HCV RNA from serum or 100 fold drop or more in HCV RNA level.</p> <p>B. For genotype 1:</p> <p>1. Undetectable HCV RNA after 24 weeks of therapy; AND</p> <p>2. One of the following:</p> <p>a. HCV RNA more than 50 IU/mL at 4 weeks into treatment</p> <p>b. Less than 100 fold drop or detectable HCV RNA 12 weeks into therapy.</p> <p>C. For genotype 3:</p> <p>1. Baseline HCV RNA more than 600,000 IU/mL; AND</p> <p>2. Steatosis or advanced fibrosis on liver biopsy.</p> <p>Hepatitis C Retreatment:</p> <p>1. One of the following</p> <p>a. Retreatment in patients who have failed or relapsed following standard or pegylated interferon monotherapy; OR</p> <p>b. For non-responders or relapsers who have significant fibrosis or cirrhosis who have undergone previous regimens of treatment using non-pegylated interferon. AND</p> <p>2. Used in combination with ribavirin.</p>				
Peg-Intron, Peg-Intron Redipen	All FDA approved indications not otherwise		<p>Hepatitis C - Treatment Naive Patients:</p> <p>1. Chronic Hepatitis C with compensated liver disease; AND</p> <p>2. Positive HCV antibody HCV RNA; AND</p>	Treatment Naive Patients: 3 years		Genotypes 5, 6: 12 weeks	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
	excluded from Part D		<p>3. HCV RNA level measurement; AND 4. Genotype test result; AND 5. For patients who have not previously been treated with interferon.</p> <p>Hepatitis C (Continuation): A. For genotypes 5 or 6: 1. Loss of detectable serum HCV RNA; OR 2. 100 fold drop or more in HCV RNA level.</p> <p>B. For genotype 1: 1. Undetectable HCV RNA after 24 weeks of therapy; AND 2. One of the following: a. HCV RNA more than 50 IU/mL at 4 weeks into treatment b. Less than 100 fold drop or detectable HCV RNA 12 weeks into therapy.</p> <p>C. For genotype 3: 1. Baseline HCV RNA more than 600,000 IU/mL; AND 2. Steatosis or advanced fibrosis on liver biopsy.</p> <p>Hepatitis C (Retreatment): 1. One of the following: a. Retreatment in patients who have failed or relapsed following standard or pegylated interferon monotherapy; OR b. For non-responders or relapsers who have significant fibrosis or cirrhosis who have undergone previous regimens of treatment using non-</p>	and older.		<p>Genotypes 2, 3: 24 weeks</p> <p>Genotypes 1, 4, co-infection with HIV/HCV: 48 weeks.</p> <p>Hepatitis C Continuation: Genotypes 1, 3: 24 weeks</p> <p>Genotypes: 5, 6: 36 weeks.</p> <p>Hepatitis C Retreatment: 1 year</p>	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			pegylated interferon. AND 2. Used in combination with ribavirin.				
Prograf intravenous	All FDA approved indications not otherwise excluded from Part D		Transplant: 1. One of the following: a. Patient received a renal (kidney), cardiac (heart), lung, pancreas, small bowel, or hepatic (liver) transplant. b. Patient received a bone marrow/stem cell transplant. AND 2. Patient is unable to take oral tacrolimus.			Length of therapy	Prograf is subject to Part B vs. Part D review (not limited to new starts only). Prograf will be approved for continuation of prior therapy if Part D.
Prograf	All FDA approved indications not otherwise excluded from Part D		Severe Uveitis: Failure to one corticosteroid. Transplant: 1. Patient received a renal (kidney), cardiac (heart), lung, pancreas, small bowel, hepatic (liver) transplant, bone marrow/stem cell transplant; AND 2. Diagnosis of graft vs. host disease in patients receiving bone marrow transplants.			Length of therapy	Prograf is subject to Part B vs. Part D review (not limited to new starts only). Prograf will be approved for continuation of prior therapy if Part D.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
Proleukin	All FDA approved indications not otherwise excluded from Part D		<p>Metastatic Renal Cell Carcinoma or Metastatic Melanoma:</p> <ol style="list-style-type: none"> 1. Measurable, histologically confirmed metastatic renal cell carcinoma or metastatic melanoma; AND 2. Good neurologic or ambulatory performance status; AND 3. Adequate organ function determined by all of the following: <ol style="list-style-type: none"> a. Normal cardiac stress test results b. FEV1 greater than 2 L on pulmonary function tests c. Creatinine concentration 1.5 mg/dL or less or calculated creatinine clearance > 60 ml/min d. Bilirubin concentration of 1.5 mg/dL or less f. SGOT/AST less than 150 IU or 4x upper limit of normal. AND 4. Platelet count greater than or equal to 100,000/ mL; AND 5. Hemoglobin greater than or equal to 10 g/dL; AND 6. WBC greater than or equal to 3,500 / mL; AND 7. At least 7 weeks since prior therapy and complete recovery from therapy-related side effects. 	All uses: 18 years and older		3 months	<p>All uses: Proleukin will be approved for continuation of prior therapy.</p> <p>Metastatic Renal Cell Carcinoma or Melanoma: Administered in a hospital setting.</p> <p>Additional treatment given only if there is some tumor shrinkage following the last course and if retreatment is not contra-indicated.</p>

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
Promacta	All FDA approved indications not otherwise excluded from Part D		<p>Chronic ITP Initial therapy:</p> <ol style="list-style-type: none"> 1. Diagnosis of ITP. 2. Platelet count less than 30,000 / microliter or less than 50,000 microliters with presence of significant mucous membrane bleeding or risk factors for bleeding. 3. Failure to corticosteroids, immunoglobulins, or splenectomy. <p>Reauthorization: Platelet count less than 50,000 / microliter</p>			<p>Initial Therapy: 6 months</p> <p>Reauthorization: 1 year</p>	
Provigil	All FDA approved indications not otherwise excluded from Part D	<p>Initial Therapy for Shift Work Sleep Disorder (SWSD): Symptoms do not meet criteria for any other sleep disorder producing insomnia or excessive sleepiness.</p>	<p>Narcolepsy: Submission of sleep study confirming the diagnosis of narcolepsy.</p> <p>Initial Therapy for Obstructive Sleep Apnea/Hypopnea Syndrome (OSAHS):</p> <ol style="list-style-type: none"> 1. Fully compliant and concurrently using continuous positive airway pressure (CPAP); AND 2. One of the following: <ol style="list-style-type: none"> a. 15 or more obstructive respiratory events; OR b. More than 5 obstructive apneas, each greater than 10 seconds in duration, per hour of sleep confirmed by a sleep study, and one of the following: <ol style="list-style-type: none"> i. Frequent arousals from sleep associated with apneas ii. Bradycardia iii. Arterial oxygen desaturation in association with apneas. iv. Unintentional sleep episodes v. Daytime sleepiness 			<p>OSAHS, SWSD: 3 months</p> <p>Other uses 12 months</p>	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<ul style="list-style-type: none"> vi. Unrefreshing sleep vii. Fatigue viii. Insomnia ix. Waking up gasping x. Loud snoring xi. Breathing interruptions during sleep <p>Reauthorization for Obstructive Sleep Apnea/Hypopnea Syndrome: Patient continues to be fully compliant on concurrent CPAP and is experiencing relief of symptomatic hypersomnolence with Provigil use.</p> <p>Shift Work Sleep Disorder:</p> <ol style="list-style-type: none"> 1. One of the following: <ul style="list-style-type: none"> a. Symptoms of excessive sleepiness or insomnia, for at least 3 months, which is temporally associated with a work period that occurs during the habitual sleep phase b. Sleep study demonstrating loss of a normal sleep-wake pattern. AND 2. Sleep disturbance causes significant distress or significant impairment; AND 3. No other disorder accounts for the symptoms. <p>Reauthorization for Shift Work Sleep Disorder:</p> <ol style="list-style-type: none"> 1. Patient is experiencing relief with use of Provigil for excessive sleepiness; AND 2. Sleep disturbance continues to cause clinically significant distress or significant impairment in occupational functioning. 				

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>Idiopathic Hypersomnia: Submission of sleep study confirming the diagnosis of Idiopathic Hypersomnia as defined by the International Classification of Sleep Disorders.</p>				
Qualaquin	All FDA approved indications not otherwise excluded from Part D	<ol style="list-style-type: none"> Severe or complicated P. falciparum malaria. Prevention of Malaria For treatment or prevention of nocturnal leg cramps. 	<p>Chloroquine-sensitive malaria: 1. Diagnosis of Malaria; AND 2. History of failure, contraindication or intolerance to chloroquine.</p> <p>Chloroquine-resistant malaria: Diagnosis of malaria.</p>			7 days	
Rebif, Rebif Titration Pack	All FDA approved indications not otherwise excluded from Part D		<p>Relapsing forms of Multiple Sclerosis: 1. Patients with relapsing forms of MS: OR 2. Patients with secondary progressive MS who continue to experience relapses.</p>			1 year	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
Reclast	All FDA approved indications not otherwise excluded from Part D		<p>Initial Therapy for Hypercalcemia of malignancy: 1. Corrected total serum calcium \geq 12 mg/dL</p> <p>Reauthorization for Hypercalcemia of malignancy: 1. Corrected total serum calcium fails to normalize after initial dose.</p> <p>Initial Therapy for Paget's disease: 1. One of the following: a. Serum alkaline phosphatase of \geq 2 times the upper limit of normal reference range b. Symptoms caused by Paget's disease c. At risk for complications</p> <p>Reauthorization for Paget's disease: 1. Serum alkaline phosphatase fails to normalize after previous therapy.</p> <p>Osteolytic bones metastases of breast cancer 1. Diagnosis of stage IV breast cancer 2. At least one lytic, metastatic bone lesion 3. Serum creatinine level below 3.0 mg/dL</p> <p>Metastatic bone lesions from solid tumors 1. Diagnosis of prostate cancer or other solid tumor 2. At least one lytic, metastatic bone lesion 3. Serum creatinine level below 3.0 mg/dL</p> <p>Osteolytic lesions of multiple myeloma 1. Diagnosis of stage III multiple myeloma 2. At least one lytic, metastatic bone lesion 3. Serum creatinine level below 3.0 mg/dL</p>			<p>Hypercalcemia, Paget's disease: One course of therapy</p> <p>Osteolytic bone metastases, metastatic bone lesions, osteolytic lesions: One year</p> <p>Osteoporosis: Length of therapy</p>	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>Prevention and treatment of osteoporosis</p> <ol style="list-style-type: none"> 1. One of the following <ol style="list-style-type: none"> a. Treatment of postmenopausal osteoporosis or osteoporosis in men, and one of the following: <ol style="list-style-type: none"> i. compression fracture in spine or peripheral fractures ii. Bone Mineral Density (BMD) T-Score ≤ -2.5 b. Prevention of postmenopausal osteoporosis in women and one of the following: <ol style="list-style-type: none"> i. BMD T-Score -1.0 to -2.5 ii. 10-year probability of a hip fracture $\geq 3\%$ or a 10-year probability of a major osteoporosis-related fracture $\geq 20\%$ based on the WHO Fracture Risk Algorithm (FRAX) c. Treatment of glucocorticoid-induced osteoporosis or prevention of glucocorticoid-induced osteoporosis in patients initiating or continuing on ≥ 7.5 mg/day oral prednisone for at least 12 months 2. One of the following: <ol style="list-style-type: none"> a. Failure to two oral bisphosphonates; OR b. Unable to tolerate oral medications 				
Regranex	All FDA approved indications not otherwise excluded from Part D		<p>Diabetic Neuropathic Ulcers</p> <ol style="list-style-type: none"> 1. Diabetic patient with ulcer wound. 2. Debridement being performed as needed; AND 2. At least two of the following are present: <ol style="list-style-type: none"> a. Stage III or IV wound b. Wound at least 1 cm x 1 cm c. Long-standing wound that does not heal with standard care d. Patients at high risk for amputation (peripheral 			Maximum: 6 months	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			neuropathy, peripheral vascular disease, skin or nail abnormalities, previous foot ulcer amputation).				
Relistor	All FDA approved indications not otherwise excluded from Part D		Initial therapy: 1. Patients with advanced illness and receiving palliative care. 2. Confirmed diagnosis of opioid-induced constipation. 3. Failure to polyethylene glycol or lactulose.	18 years and older		6 months	
Remicade	All FDA approved indications not otherwise excluded from Part D		Rheumatoid Arthritis: 1. Diagnosis of moderate-to-severe RA 2. Concurrently on methotrexate or failure to methotrexate or 2 DMARDs (azathioprine, cyclosporine, gold, hydroxychloroquine, leflunomide, penicillamine, sulfasalazine) for 3 months. Psoriatic Arthritis: 1. Diagnosis of active Psoriatic arthritis 2. Failure or contraindication to methotrexate or 2 of the following for 3 months: cyclosporine, gold, leflunomide, or sulfasalazine for 3 months. Ankylosing Spondylitis (AS): 1. Diagnosis of AS 2. Failed 2 NSAIDs for 3 months Plaque Psoriasis: 1. Diagnosis of moderate-to-severe (greater than 6 months) plaque psoriasis; AND 2. Failure, contraindication, intolerance or unavailability	RA, PsA, AS, Plaque Psoriasis, FCD, UC: 18 years and older. Crohn's Disease: 6 years and older.	RA, AS, PsA: Prescribed or recommended by a rheumatologist. Crohn's Disease, Fistulizing Crohn's Disease, UC: Prescribed by a gastroenterologist or by gastroenterologist consult. Plaque Psoriasis:	12 months	Verification that the patient has been evaluated for tuberculosis (TB) and treated accordingly.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>of phototherapy and one of the following: methotrexate, cyclosporine, acitretin, hydroxyurea, sulfasalazine, 6-thioguanine, mycophenolate.</p> <p>Crohn's Disease (CD): 1. Moderate to severe CD; AND 2. Failed one of the following: corticosteroids, 6-mercaptopurine, azathioprine, methotrexate, aminosalicylate</p> <p>Fistulizing Crohn's Disease (FCD): 1. Draining fistulas for 3 months; AND 2. On or failed one of the following: 6-mercaptopurine, azathioprine, antibiotics, oral corticosteroids, methotrexate</p> <p>Ulcerative Colitis (UC): 1. Moderate to severe UC; AND 2. Failed on one of the following: corticosteroids, 5-aminosalicylic acid, azathioprine, 6-mercaptopurine, cyclosporine</p> <p>Sarcoidosis: Failed one steroid and one immunosuppressant.</p> <p>Reauthorization: Demonstration of clinical response to therapy</p>		<p>Prescribed or recommended by a dermatologist.</p> <p>Sarcoidosis: Prescribed or recommended by a pulmonologist.</p>		
Remodulin	All FDA approved indications not		<p>Pulmonary Arterial Hypertension Patients with a confirmed diagnosis of pulmonary arterial hypertension (modified WHO Group I) which is</p>			Length of therapy	Remodulin is subject to Part B vs. Part D

Unison Advantage® (HMO) products are offered by UnitedHealthcare Insurance Company and its affiliated companies, Medicare Advantage organizations with a Medicare contract.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
	otherwise excluded from Part D		symptomatic.				review.
Revatio	All FDA approved indications not otherwise excluded from Part D	Pulmonary Arterial Hypertension (PAH): Patients using organic nitrates.	Pulmonary Arterial Hypertension: Patients with a confirmed diagnosis of pulmonary arterial hypertension (modified WHO Group I) which is symptomatic.			Length of therapy	
Revlimid	All FDA approved indications not otherwise excluded from Part D		Myelodysplastic Syndrome (MDS): 1. Diagnosis of myelodysplastic syndrome associated with a deletion 5q cytogenic abnormality; AND 2. Patient is transfusion dependent. OR 1. Diagnosis of myelodysplastic syndrome without a deletion 5q cytogenic abnormality; AND 2. Failure of initial treatment with epoetin alfa or darbopoetin alfa, hypomethylating agents (e.g., Vidaza, Dacogen), or immunosuppressive therapy (e.g., antithymocyte globulin, cyclosporine). Multiple Myeloma: Used in combination with dexamethasone. Chronic Lymphocytic Leukemia (CLL): Relapsed or refractory to one prior therapy for CLL.		MDS, Multiple Myeloma, CLL: Prescribed by an oncologist or hematologist or by oncology or hematology consult.	MDS, Multiple Myeloma: 6 months	Revlimid will be approved for continuation of prior therapy.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
Rebetol, Ribasphere, ribavirin	All FDA approved indications not otherwise excluded from Part D		Hepatitis C: Adults with a diagnosis of Hepatitis C with compensated liver disease, and verification of concurrent use with an alfa-interferon product.			Length of therapy	
Rituxan	All FDA approved indications not otherwise excluded from Part D Chronic Lymphocytic Leukemia Immune or idiopathic thrombocytopenic purpura Waldenstrom's macroglobulinemia		Non-Hodgkin's Lymphoma: One of the following: 1. As first-line treatment of diffuse large B-cell, CD20-positive, non-Hodgkin's lymphoma in combination with CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone) or other anthracycline-based chemotherapy regimens 2. As first-line treatment of follicular, CD20-positive, B-cell non-Hodgkin's lymphoma in combination with CVP (cyclophosphamide, vincristine, prednisolone/prednisone) chemotherapy 3. For the treatment of low-grade, CD20-positive, B-cell non-Hodgkin's lymphoma in patients with stable disease or who achieve a partial or complete response following first-line treatment with CVP chemotherapy 4. Confirmed diagnosis of relapsed or refractory, low grade or follicular CD20-positive, B-cell non-Hodgkin's lymphoma. Initial Therapy for Rheumatoid Arthritis (RA): 1. Diagnosis of moderate/severe RA; AND 2. Used in combination with or failure to methotrexate;	RA: 18 years and older.	RA: Prescribed by a rheumatologist.	All uses except RA: 1 year RA: 1 month	

Unison Advantage® (HMO) products are offered by UnitedHealthcare Insurance Company and its affiliated companies, Medicare Advantage organizations with a Medicare contract.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>AND</p> <p>3. Failure to a TNF antagonist.</p> <p>Reauthorization for Rheumatoid Arthritis:</p> <p>1. Documented positive clinical response; AND</p> <p>2. At least 16 weeks since last Rituxan treatment.</p>				
Sabril	All FDA approved indications not otherwise excluded from Part D		<p>Complex Partial Seizures (CPS):</p> <p>For use as adjunctive therapy in patients who have failed two formulary anticonvulsants.</p> <p>Infantile Spasms (IS):</p> <p>Diagnosis of IS.</p>			12 months	Approve for continuation of prior therapy.
Samsca	All FDA approved indications not otherwise excluded from Part D		<p>Initial therapy for hypervolemic and euvolemic hyponatremia:</p> <p>1. Diagnosis of significant hyponatremia (euvolemic or hypervolemic); AND</p> <p>2. Treatment has been initiated or re-initiated in a hospital setting prior to discharge.</p> <p>Reauthorization for hypervolemic and euvolemic hyponatremia:</p> <p>1. Documentation of clinical benefit; AND</p> <p>2. Treatment has been initiated or re-initiated in a hospital setting prior to discharge.</p>		Prescribed by an endocrinologist or nephrologist or by consultation with an endocrinologist or nephrologist	<p>Initial: 1 month.</p> <p>Reauth: 3 months</p>	
octreotide acetate	All FDA approved indications not		<p>Acromegaly:</p> <p>1. Inadequate response to surgery and/or radiotherapy or patients who are not a surgical and/or radiotherapy</p>			Acromegaly: Long term approval.	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
	otherwise excluded from Part D		<p>candidate</p> <p>2. Diagnosis of acromegaly by one of the following:</p> <ul style="list-style-type: none"> a. Serum growth hormone (GH) level greater than 1 ng/mL after a 2-hour oral glucose tolerance test b. Elevated serum IGF-1 levels as compared to normal reference values by age and gender. <p>Carcinoid Tumors: Diagnosis of metastatic carcinoid tumor for symptomatic treatment of severe diarrhea or flushing.</p> <p>Vasoactive Intestinal Peptide Tumors: Diagnosis of metastatic vasoactive peptide tumor, for symptomatic treatment of diarrhea associated with vasoactive peptide tumor.</p> <p>Cancer Chemotherapy Induced Diarrhea:</p> <ul style="list-style-type: none"> 1. Diagnosis of diarrhea due to concurrent cancer chemotherapy; OR 2. Both of the following: <ul style="list-style-type: none"> a. Diagnosis of complicated diarrhea due to concurrent cancer chemotherapy b. History of failure to standard therapy. <p>AIDS-related Diarrhea:</p> <ul style="list-style-type: none"> 1. Diagnosis of AIDS-related diarrhea. 2. History of failure to standard therapy 			<p>Tumors: 6 months</p> <p>Chemotherapy induced diarrhea, AIDS-related diarrhea: 3 months</p>	
Sandostatin LAR Depot	All FDA approved indications not		<p>Acromegaly:</p> <ul style="list-style-type: none"> 1. Inadequate response to surgery and/or radiotherapy or patients who are not a surgical and/or radiotherapy 			<p>Acromegaly: Long term approval.</p>	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
	otherwise excluded from Part D		<p>candidate.</p> <p>2. Patient has shown to respond to and tolerate octreotide injection for at least 2 weeks.</p> <p>Carcinoid Tumors:</p> <p>1. Diagnosis of metastatic carcinoid tumor, for symptomatic treatment of severe diarrhea or flushing; AND</p> <p>2. Patient has been shown to respond to and tolerate octreotide.</p> <p>Vasoactive Intestinal Peptide Tumors:</p> <p>1. Diagnosis of metastatic vasoactive peptide tumor, for symptomatic treatment of diarrhea associated with vasoactive peptide tumor; AND</p> <p>2. Patient has been shown to respond to and tolerate octreotide.</p>			Tumors: 6 months	
Saphris	All FDA approved indications not otherwise excluded from Part D		Failure on one tier 1 or tier 2 atypical antipsychotic			12 months	
Serostim	All FDA-approved indications not otherwise excluded from Part D		<p>Initial Therapy for AIDS associated cachexia or wasting:</p> <p>1. Documented AIDS-associated cachexia or wasting defined as one of the following:</p> <p>a. 10% unintentional weight loss over 12 months</p> <p>b. 7.5% unintentional weight loss over 6 months</p>			AIDS wasting: 3 months	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>c. 5% body cell mass (BCM) loss within 6 months, or body mass index (BMI) less than 20 kg/m²</p> <p>d. In men, BCM less than 35% of TBW and BMI less than 27 kg/m²</p> <p>e. In women, BCM less than 23% of TBW and BMI less than 27 kg/m²</p> <p>. AND</p> <p>2. Nutritional evaluation since onset of wasting first occurred.</p> <p>3. Other causes of HIV-associated wasting have been ruled out.</p> <p>4. Target or goal weight, BCM, or BMI.</p> <p>3. For male patients: Screen for hypogonadism (obtain free testosterone levels)</p> <p>4. For male patients: Failure to respond to testosterone replacement therapy in patients with hypogonadism.</p> <p>Reauthorization for AIDS associated cachexia or wasting:</p> <p>1. Positive response to therapy based on greater than or equal to 2% increase in body weight and/or BCM.</p> <p>2. Target or goal weight, BCM, BMI has not been achieved.</p> <p>Reinitiation of therapy for AIDS associated cachexia or wasting:</p> <p>1. Positive response to therapy based on greater than or equal to 2% increase in body weight and/or BCM in a previous treatment.</p> <p>2. Evidence of HIV-associated cachexia or wasting (as defined above) after an 8-week observation period.</p>				

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
Simponi	All FDA-approved indications not otherwise excluded from Part D		<p>Rheumatoid Arthritis (RA):</p> <ol style="list-style-type: none"> Diagnosis of moderate-to-severe RA; AND Concurrently on methotrexate or failed methotrexate or 2 DMARDs (azathioprine, cyclosporine, gold, hydroxychloroquine, leflunomide, penicillamine, sulfasalazine) for 3 months. <p>Psoriatic Arthritis (PsA):</p> <ol style="list-style-type: none"> Diagnosis of active PsA. Failure or contraindication to methotrexate or 2 of the following for 3 months: cyclosporine, gold, leflunomide, or sulfasalazine. <p>Ankylosing Spondylitis (AS):</p> <ol style="list-style-type: none"> Diagnosis of AS. Failed 2 NSAIDs for 3 months. <p>Reauthorization (all diagnoses): Demonstration of clinical response to therapy.</p>	<p>RA, PsA, AS: 18 years and older</p>	<p>RA (Initial) or AS (Initial): Prescribed or recommended by a rheumatologist.</p> <p>PsA: Prescribed or recommended by a rheumatologist or dermatologist.</p>	<p>Initial Therapy: 12 months.</p> <p>Reauthorization (all uses): 12 months.</p>	All diagnoses require verification that the patient has been evaluated for tuberculosis and has been treated accordingly
Somatuline Depot	All FDA-approved indications not otherwise excluded from Part D		<p>Acromegaly:</p> <ol style="list-style-type: none"> Patients who require long-term treatment due to: <ol style="list-style-type: none"> Inadequate response to surgery and/or radiotherapy; OR Who are not surgical and/or radiotherapy candidates. AND Diagnosis of acromegaly by one of the following: <ol style="list-style-type: none"> Serum growth hormone level greater than 1 ng/mL after a 2-hour oral glucose tolerance test; OR Elevated serum IGF-1 levels as compared to 			Indefinite, long-term therapy (open-ended)	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			normal reference values by age and gender.				
Somavert	All FDA-approved indications not otherwise excluded from Part D		<p>Initial Therapy for Acromegaly:</p> <ol style="list-style-type: none"> 1. One of the following: <ol style="list-style-type: none"> a. Inadequate response to surgery and/or radiation therapy b. Not a candidate for surgery or radiation. AND 2. Inadequate response or intolerance to octreotide, or lanreotide, or IGF-1 value greater than 900 ng/mL. <p>Reauthorization for Acromegaly: Serum IGF-1 level within the age-adjusted normal range.</p>			<p>Acromegaly (initial): 12 weeks</p> <p>Acromegaly (reauthorization): Indefinite</p>	
Sporanox solution	All FDA approved indications not otherwise excluded from Part D		<p>Fungal Infection:</p> <ol style="list-style-type: none"> 1. Diagnosis of one of the following: <ol style="list-style-type: none"> a. Blastomycosis b. Histoplasmosis c. Aspergillosis 2. Onychomycosis in patients unable to swallow tablets; OR 3. Diagnosis of febrile neutropenia with suspected fungal infection, or oropharyngeal or esophageal candidiasis. 			Length of therapy	
Sprycel	All FDA approved indications not otherwise excluded from Part D		<p>Chronic Myeloid Leukemia (CML):</p> <ol style="list-style-type: none"> 1. Diagnosis of Philadelphia chromosome positive or BCR-ABL positive chronic, accelerated, or myeloid or lymphoid blast phase chronic myeloid leukemia; AND 2. Failure to Gleevec. <p>Acute Lymphoblastic Leukemia (ALL):</p>			Length of therapy	Sprycel will be approved for continuation of prior therapy.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<ol style="list-style-type: none"> 1. Diagnosis of Philadelphia chromosome-positive acute lymphoblastic leukemia; AND 2. Failure to Gleevec. 				
Stelara	All FDA approved indications not otherwise excluded from Part D		<p>Plaque Psoriasis</p> <ol style="list-style-type: none"> 1. Diagnosis moderate-to-severe plaque psoriasis. 2. Failed phototherapy and systemic therapy with one of the following: methotrexate, cyclosporine, acitretin, hydroxyurea, sulfasalazine, 6-thioguanine, or mycophenolate. 3. Failed one TNF blocker. 4. Verification that the patient has been evaluated for TB and has been treated accordingly. <p>Reauthorization: Demonstration of clinical response to therapy.</p>	Age greater than or equal to 18 years.	Prescribed or recommended by a dermatologist.	12 amonths	Authorization will be issued for 45 mg (for patients weighing less than or equal to 100 kg) or 90 mg (for patients weighing more than 100 kg) initially and 4 weeks later followed by 45 mg (for patients weighing less than or equal to 100 kg) or 90 mg (for patients weighing more than 100 kg) every 12 weeks for 12 months.
Sutent	All FDA		Gastrointestinal Stromal Tumor (GIST):		Prescribed by	12 months	Sutent will be

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
	approved indications not otherwise excluded from Part D		Disease progression on or intolerance to Gleevec. Renal Cell Carcinoma: One of the following: 1. Diagnosis of renal cell carcinoma with relapse following surgical excision 2. Diagnosis of renal cell carcinoma with medically or surgically unresectable tumor 3. Diagnosis of Stage IV renal cell carcinoma.		an oncologist.		approved for continuation of prior therapy.
Symlin, Symlinpen	All FDA approved indications not otherwise excluded from Part D		Diabetes Mellitus: 1. Type 1 or type 2 diabetes 2. Concurrent use of insulin therapy	18 years and older.		Length of therapy	
Tarceva	All FDA approved indications not otherwise excluded from Part D		Non-Small Cell Lung Cancer (NSCLC): Patients diagnosed with locally advanced or metastatic NSCLC after failure of at least one prior chemotherapy regimen. Pancreatic Cancer: 1. Patient diagnosed with locally advanced, unresectable or metastatic pancreatic cancer. 2. Used in combination with gemcitabine. Reauthorization (all uses): Patient has not experienced disease progression.		Prescribed by an oncologist.	6 months	Tarceva will be approved for continuation of prior therapy.
Targretin (oral)	All FDA approved		Definitive diagnosis of cutaneous T-cell lymphoma (CTCL)			12 months	Targretin will be approved

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
	indications not otherwise excluded from Part D						for continuation of prior therapy.
Tasigna	All FDA approved indications not otherwise excluded from Part D		Chronic Myelogenous Leukemia: 1. Diagnosis of Philadelphia chromosome positive chronic or accelerated phase chronic myeloid leukemia; AND 2. Failure to Gleevec.			Length of therapy	Tasigna will be approved for continuation of prior therapy.
Testosterone injectable	All FDA-approved indications not otherwise excluded from Part D.		Hypogonadism: Diagnosis of male hypogonadism with a pre-treatment total testosterone level below normal physiological value (less than 280 ng/dl), or pre-treatment free testosterone below normal reference value. Delayed puberty: Diagnosis of delayed puberty in males.			Hypogonadism: Long-term. Delayed puberty: 6 months	
Topical testosterone: Androderm, Androgel	All FDA-approved indications not otherwise excluded from Part D.		Hypogonadism: Diagnosis of hypogonadism in men with a pre-treatment testosterone level below normal physiological value of 280 ng/dL or below normal reference level provided by the physician laboratory.			Length of therapy	
Thalomid	All FDA-approved indications not		Erythema Nodosum Leprosum (ENL): Confirmed diagnosis of moderate to severe ENL.			AS: 1 month ENL, MM: 1	Thalomid will be approved for

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
	<p>otherwise excluded from Part D.</p> <p>Waldenstrom's Macro-globulinemia (WM)</p> <p>Aphthous stomatitis or ulcers (AS)</p> <p>Crohn's Disease,</p> <p>Graft-versus-Host Disease (GVHD)</p> <p>Primary Brain Tumors</p> <p>AIDS-related cachexia or wasting</p> <p>Renal Cell Carcinoma</p>		<p>Multiple Myeloma (MM): Confirmed diagnosis of MM.</p> <p>Waldenstrom's Macro-globulinemia (WM): Confirmed diagnosis of WM.</p> <p>Aphthous stomatitis (AS) or ulcers: 1. One of the following: a. Diagnosis of HIV-associated aphthous ulcers b. Recurrent aphthous stomatitis in immunocompromised patients. AND 2. Failure to alternative therapies.</p> <p>Crohn's Disease: Failure to all of the following standard treatment regimens: 1. Corticosteroids 2. 5-aminosalicylic acid 3. Immunomodulators 4. Remicade.</p> <p>Graft-versus-Host Disease (GVHD): 1. Diagnosis of chronic or refractory GVHD; AND 2. Failure to corticosteroids or one of the following: a. Azathioprine b. Tacrolimus c. Cyclosporine d. Antithymocyte globulin e. mycophenolate</p> <p>Primary Brain Tumors:</p>			<p>year</p> <p>WM, GVHD, and Primary Brain Tumors, RCC: 6 months</p> <p>Other Uses: 3 months</p>	<p>continuation of prior therapy.</p>

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
			<p>1. As adjuvant therapy to current cytotoxic therapies; OR 2. Previous failure to cytotoxic therapies and/or tumor resection.</p> <p>Initial Therapy for AIDS-related cachexia or wasting: 1. Diagnosis of AIDS wasting or cachexia; AND 4. Failure, contraindication or intolerance to standard treatments.</p> <p>Advanced Renal Cell Carcinoma: 1. Confirmed diagnosis of metastatic renal cell carcinoma; AND 2. Failure to two of the following: a. Interferon-alfa-2b b. Interleukin-2 c. Sorafenib d. Sunitanib e. temsirolimus f. pazopanib</p>				

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
trimethoben z-amide	All FDA approved indications not otherwise excluded from Part D			No prior authorization if patient is less than 65 years old		Length of therapy	
Topical Retinoids: Avita, Retin A Micro, Tretin-X, tretinoin	All FDA approved indications not otherwise excluded from Part D					12 months	
Tracleer	All FDA approved indications not otherwise excluded from Part D		Pulmonary Arterial Hypertension: Patients with a confirmed diagnosis of pulmonary arterial hypertension (modified WHO Group I) which is symptomatic.			Length of therapy	
Treanda	All FDA approved		Non-Hodgkin's Lymphoma (NHL): 1. Diagnosis of indolent B-cell NHL.			6 months	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
	indications not otherwise excluded from Part D		2. Progression of NHL during or within 6 months of treatment with rituximab or a rituximab-containing regimen.				
Tykerb	All FDA approved indications not otherwise excluded from Part D		Breast Cancer: 1. Diagnosis of HER2-positive advanced or metastatic breast cancer 2. Confirmation of normal left ventricular ejection fraction.			Length of therapy	Tykerb will be approved for continuation of prior therapy.
Tysabri	All FDA approved indications not otherwise excluded from Part D		Relapsing forms of Multiple Sclerosis (MS): Failure to one of the following: Avonex, Betaseron, Copaxone, Rebif. Initial Therapy for Crohn's Disease (CD): 1. Moderate-to-severe Crohn's disease with evidence of inflammation; AND 2. History of conventional therapy. History of a TNF blocker. Patient is not receiving immunosuppressants. Reauthorization for Crohn's Disease: Demonstrated remission or significant clinical response to Tysabri.	Initial therapy for CD: 18 years and older.	Relapsing MS: Prescribing physician is enrolled in the TOUCH Prescribing Program. Initial Therapy - CD: Prescribing physician is enrolled in the CD TOUCH Prescribing Program.	Initial Therapy: Tysabri will be authorized for 1 year for MS and 3 months for CD Reauthorization for CD: Tysabri will be reauthorized for 6 months for patients on steroids. Otherwise, 3 months.	For Relapsing MS: Tysabri will not be authorized in combination with Avonex, Betaseron, Copaxone, or Rebif.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
Vancocin	All FDA approved indications not otherwise excluded from Part D		Pseudo-membranous Colitis: 1. Diagnosis of pseudo-membranous colitis due to Clostridium difficile; AND 2. Failure to oral Flagyl.			Length of therapy	
Vectibix	All FDA approved indications not otherwise excluded from Part D		Colorectal Cancer: 1. Diagnosis of metastatic colorectal cancer. 2. Relapsed, refractory, or disease progression on one standard chemotherapy regimen containing a fluoropyrimidine, oxaliplatin, or irinotecan. 3. Tumor expresses wild-type KRAS gene.			6 months	Vectibix will be approved for continuation of prior therapy.
Ventavis	All FDA approved indications not otherwise excluded from Part D		Pulmonary Arterial Hypertension Patients with a confirmed diagnosis of pulmonary arterial hypertension (modified WHO Group I) which is symptomatic.			Length of therapy	Ventavis is subject to Part B vs. Part D review.
Vimpat	All FDA-approved indications not otherwise excluded from Part D.					Length of therapy	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
Vpriv	All FDA-approved indications not otherwise excluded from Part D		Gaucher's disease 1. Confirmed diagnosis of Type 1 Gaucher's disease. 2. Symptomatic disease defined by one of the following: a. moderate to severe anemia b. thrombocytopenia c. bone disease d. hepatomegaly e. splenomegaly.			12 months	
Votrient	All FDA-approved indications not otherwise excluded from Part D		Renal cell carcinoma: Diagnosis of advanced renal cell cancer.			1 year	Prior authorization applies to new starts only

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
Xenazine	All FDA approved indications not otherwise excluded from Part D. Tardive dyskinesia. Tourette's syndrome.		<p>Initial therapy for Huntington's Disease: Diagnosis of chorea in patients with Huntington's disease.</p> <p>Initial therapy for Tardive dyskinesia and Tourette's syndrome: 1. Stereotypes associated with tardive dyskinesia; OR 2. Both of the following: a. Tics associated with Tourette's syndrome; AND b. Failure, contraindication, or intolerance to Haldol (haloperidol)</p> <p>Reauthorization for Huntington's Disease, Tardive dyskinesia, and Tourette's syndrome: Documented clinical response and benefit from therapy.</p>		Prescribed by a neurologist.	<p>Initial Therapy: 3 months.</p> <p>Reauthorization: 12 months.</p>	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
Xolair	All FDA approved indications not otherwise excluded from Part D		<p>Initial Therapy for Allergic Asthma:</p> <ol style="list-style-type: none"> 1. Diagnosis of moderate-to-severe persistent allergic asthma, defined by one of the following: <ol style="list-style-type: none"> a. Daily asthmatic symptoms b. Daily use of inhaled short-acting beta agonists c. Exacerbations affect/limit activity d. Exacerbations 2 or more times per week e. Nocturnal symptoms once a week or more f. Forced expiratory volume in one second or peak expiratory flow less than or equal to 80% of predicted g. PEF variability greater than 30%. AND 2. Baseline IgE level greater than or equal to 30 IU/mL; AND 3. Documented failure to combination therapy with an inhaled corticosteroid at the maximum dosage and a long-acting beta-agonist. <p>Reauthorization for Allergic Asthma:</p> <ol style="list-style-type: none"> 1. Documented reduction in the frequency of asthma exacerbations while treated with Xolair; AND 2. Documented reduction in the use of rescue medications or inhaled corticosteroids while treated with Xolair. 	Initial treatment: 6 years and older.	Initial Therapy: Prescribed by a pulmonologist or allergist/immunologist.	Initial Therapy: 16 weeks Reauthorization: 1 year	

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
Zolinza	All FDA approved indications not otherwise excluded from Part D		Definitive diagnosis of cutaneous T-cell lymphoma (CTCL)			12 months	Zolinza will be approved for continuation of prior therapy.
Zorbtive	All FDA approved indications not otherwise excluded from Part D		Short Bowel Syndrome (SBS): 1. All of the following: a. Diagnosis of SBS. b. BMI 17 to 28 kg/m2. d. At least 2 months post-resection. e. Patient is currently on parenteral nutrition f. Patient is able to ingest some food g. Intact stomach and duodenum			4 weeks	
Zortress	All FDA-approved indications not otherwise excluded from Part D. Cardiac Transplant.		Transplant: Patient received a renal or cardiac transplant.			12 months	Subject to Part B vs. Part D review. Approve for continuation of prior therapy if Part D.

Unison Advantage® (HMO)



Non-Dual Plans:

Unison Advantage® Basic (HMO)

Unison Advantage® Choice (HMO)

Drugs	Covered Uses	Exclusion Criteria	Required Medical Information	Age Restrictions	Prescriber Restrictions	Coverage Duration	Other Criteria
Zyvox	All FDA approved indications not otherwise excluded from Part D		<p>Infections: One of the following:</p> <ol style="list-style-type: none"> 1. Infections caused by vancomycin-resistant enterococci (VRE) documented by culture and sensitivity report. 2. Nosocomial pneumonia caused by methicillin-resistant Staphylococcus aureus (MRSA) documented by culture and sensitivity report 3. Complicated skin and skin structure infections (including diabetic foot infections) without osteomyelitis caused by MRSA documented by culture and sensitivity report. 4. Empirical treatment of patients with community-acquired complicated skin and skin structure infections without osteomyelitis where MRSA infection is likely, in patients who have failed one of the following: <ol style="list-style-type: none"> a. trimethoprim-sulfamethoxazole b. tetracycline c. doxycycline d. minocycline. 5. Chronic osteomyelitis or prosthetic joint infection due to MRSA or methicillin-resistant Staphylococcus epidermidis (MRSE), and failure to two or resistance to all of the following: <ol style="list-style-type: none"> a. trimethoprim-sulfamethoxazole b. fluoroquinolone c. tetracycline d. clindamycin 6. Cutaneous or systemic infection due to Nocardia, and failure to two or resistance to all of the following: <ol style="list-style-type: none"> a. trimethoprim-sulfamethoxazole b. third-generation cephalosporin c. imipenem-cilastatin d. amikacin e. minocycline f. amoxicillin-clavulanate 7. As continuation of therapy when transitioning from intravenous daptomycin, intravenous vancomycin, or intravenous Zyvox therapy. 		<p>Chronic osteomyelitis, prosthetic joint infection, cutaneous or systemic Nocardiosis: Prescribed by an infectious disease specialist or by consultation with an infectious disease specialist.</p>	<p>Chronic osteomyelitis, prosthetic joint Infection: 6 weeks</p> <p>Cutaneous Nocardiosis: 3 months</p> <p>Systemic Nocardiosis: 12 months</p> <p>Other uses: 28 days</p>	

Unison Advantage® (HMO) products are offered by UnitedHealthcare Insurance Company and its affiliated companies, Medicare Advantage organizations with a Medicare contract.

Unison Advantage® (HMO)



Non-Dual Plans:

- Unison Advantage® Basic (HMO)
- Unison Advantage® Choice (HMO)

The following drugs may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination:

- | | | |
|---------------------------------|-----------------------------------------|---------------------------------------|
| • Abelcet | • Doxil | • Ondansetron ODT |
| • Acetylcysteine | • Doxorubicin | • Orthoclone |
| • Acyclovir | • Engerix-B | • Premasol |
| • Adriamycin | • Foscarnet | • Procalamine |
| • Albuterol | • Freamine HBC | • Prosol |
| • Ambisome | • Freamine III | • Pulmicort |
| • Aminess | • Gengraf | • Pulmozyme |
| • Aminosyn (all except 15% INJ) | • Granisetron | • Rapamune |
| • Amphotec | • Granisol | • Recombivax HB |
| • Amphotericin | • Hepatamine | • Renamin |
| • Atgam | • Hepatasol | • Sandimmune (all except 50mg/ml INJ) |
| • Clinimix | • Intralipid | • Simulect |
| • Clinimix E | • Ipratropium bromide | • Thymoglobulin |
| • Clinisol SF | • Ipratropium bromide/albuterol sulfate | • Tobi |
| • Cromolyn Sodium | • Myfortic | • Travasol |
| • Cyclophosphamide | • Nebupent | • Trophamine (all except 6% INJ) |
| • Cyclosporine | • Nephramine | • Zenapax |
| • Cyclosporine Modified | • Novamine | • Zuplenz |
| • Cytovene | • Ondansetron | |